

Letter to the Editor: "Secondary Cancers After Ewing Sarcoma and Ewing Sarcoma as Second Malignant Neoplasm"

The development of a second malignant neoplasm (SMN) is the most devastating event after the successful treatment of childhood cancer. Several studies have estimated an increased (22–35%) cumulative risk of secondary cancers following Ewing sarcoma [1,2]. Radiation-induced osteosarcoma and therapy-related acute myeloid leukemia (AML) are the most frequent SMNs, although malignant fibrous histiocytoma, breast carcinoma, and other common tumours in adults have been described [1–7]. In a recent review from three institutions, 16 cases of SMN were reported among 266 survivors of Ewing sarcoma [3]. Most of these tumours developed within previously irradiated fields, and a radiation dose-dependency was found to be evident for secondary sarcomas. In contrast, Ewing sarcoma has rarely been observed as a SMN, with only about 15 such cases reported [8]. While most of these occurrences were associated with retinoblastoma [9], isolated cases have been reported following treatment of other tumours like large cell lymphoma or Wilms tumour [8,10]. No cases of Ewing sarcoma were found among 91 secondary bone sarcomas from the Late Effects Study Group [11].

A retrospective review was undertaken of 121 patients with Ewing sarcoma of bone or soft tissue and primitive neuroectodermal tumours seen in our single centre from 1970 through 1995. The purpose was to ascertain the incidence of second cancers. Primary treatment consisted of systemic multiagent chemotherapy plus local irradiation, wide surgery, or both. Forty-two (35%) patients were alive and disease-free more than 3 years after diagnosis. Table I summarizes the characteristics of 4 patients found to have a SMN after the treatment of Ewing sarcoma. Case 3 has been previously reported [12]. One breast carcinoma could be related to local irradiation (whole lung radiation therapy for metastatic disease), whereas both acute leukemias were probably chemotherapy-related. Interestingly, only one case of osteosarcoma was seen occurring in a patient surgically treated for Ewing sarcoma. Moreover, in three other patients Ewing sarcoma developed as a second cancer (Table II). None of them could be related to radiation therapy of the primary tumour. Aside from case 2, who received symptomatic measures, patients were treated with standard

protocols and four of them are now alive and disease-free.

Data from the literature indicate that patients with Ewing sarcoma are at a significantly increased risk of SMNs [1,4]. Therapy-related factors (radiotherapy, alkylating agents, and epipodophyllotoxins) have been most often advocated. However, some observations also suggest a genetic tumour predisposition in this population: (1) There is an excess risk of SMNs following Ewing sarcoma in comparison with other childhood cancers [4]; (2) The incidence of Ewing sarcoma as a SMN could be higher than suspected, since such cases are usually excluded from clinical trials; (3) SMNs, including osteosarcoma, have been described in survivors of Ewing sarcoma, which developed outside the irradiation fields of the primary tumour; (4) An inherent relationship between Ewing sarcoma and AML has been suggested, possibly in terms of a common histogenesis [5,13]. Although numbers are small and unrelated coincidence can not be excluded, our experience suggests that there is an intrinsic predisposition to SMNs in patients with Ewing sarcoma and that it is not only a therapy-related matter.

Jorge Aparicio, MD
Angel Segura, MD
Joaquín Montalar, MD
Department of Medical Oncology
Hospital La Fe
Valencia, Spain

Amparo Verdeguer, MD
Victoria Castel, MD
Department of Paediatric Oncology
Hospital La Fe
Valencia, Spain

A. Beatriz Sánchez-Heras, MD
Department of Medical Oncology
Hospital San Juan
Alicante, Spain

TABLE I. Second Malignant Neoplasms Among 121 Patients With Ewing Sarcoma*

| Patient no. | Ewing sarcoma | | | | Second cancer | |
|-------------|---------------|-------------------|----------------------------------------------------------------------|-------------------------------|--------------------|-----------------------------------|
| | Age/sex | Site Stage | Treatment | Interval (years) ^a | Histology Site | Current status (months after SMN) |
| 1 | 11/F | Radius Metastatic | VCR-ADR RT 50 Gy, lung 20 Gy | 15 | Ductal Ca. Breast | Died of SMN 52 |
| 2 | 6/M | Ribs Metastatic | SUR + RT 30 Gy VCR, ADR, CPA, DACT, CDDP, IFO, DTIC, VP-16, VM-26 | 9 | AML (M2) | Died of SMN 3 |
| 3 | 8/M | Pelvis Localized | SUR CPA, ADR, IFO, VP-16, VCR, DACT, CDDP, VM-26 | 2.6 | AML (M3) | Alive, disease-free 48+ |
| 4 | 10/M | Pelvis Localized | SUR CPA, ADR, VCR, DACT, BCNU | 10 | Osteosarcoma Femur | Alive, disease-free 9+ |

*F: female; M: male; RT: radiation therapy; SUR: surgery; Ca: carcinoma; SMN: second malignant neoplasm; AML: acute myeloid leukemia; VCR: vincristine; ADR: doxorubicin; CPA: cyclophosphamide; DACT: dactinomycin; CDDP: cisplatin; IFO: ifosfamide; DTIC: dacarbazine; VP-16: etoposide; VM-26: teniposide; BCNU: carmustine.

^aInterval from diagnosis of primary tumor to diagnosis of second malignancy.

TABLE II. Ewing Sarcoma as Second Malignant Neoplasm*

| Patient no. | Primary tumor | | | Interval (years) ^a | Ewing sarcoma | |
|-------------|---------------|----------------|--------------------------------------------|-------------------------------|------------------|-----------------------------------|
| | Age/sex | Histology Site | Treatment | | Site Stage | Current status (months after SMN) |
| 5 | 25/M | MFH Scapula | SUR + RT 60 Gy CYVADIC | 5.5 | Pelvis Localized | Died of SMN 30 |
| 6 | 3/M | ALL | VCR, LAG, ADR, 6-MP, MTX, AraC, Cranial RT | 12 | Ulna Metastatic | Alive, disease-free 15+ |
| 7 | 11/M | AML (M4) | ADR, DNR, MTZ, AraC, 6-TG, 6-MP, VP-16 | 7 | Fibula Localized | Alive, disease-free 20+ |

*M: male; MFH: malignant fibrous histiocytoma; RT: radiation therapy; SUR: surgery; SMN: second malignant neoplasm; CYVADIC: cyclophosphamide, vincristine, doxorubicin, and dacarbazine; AML: acute myeloid leukemia; ALL: acute lymphoblastic leukemia; VCR: vincristine; LAG: L-asparaginase; ADR: doxorubicin; AraC: cytarabine; 6-MP: 6-mercaptopurine; MTX: methotrexate; DNR: daunorubicin; MTZ: mitoxantrone; 6-TG: thioguanine; VP-16: etoposide.

^aInterval from diagnosis of primary tumor to diagnosis of second malignancy.

REFERENCES

- Strong LC, Herson J, Osborne BM, et al.: Risk of radiation-related subsequent malignant tumors in survivors of Ewing's sarcoma. *J Natl Cancer Inst* 62:1401-1406, 1979.
- Tucker MA, D'Angio GJ, Boice JD Jr, et al.: Bone sarcomas linked to radiotherapy and chemotherapy in children. *N Engl J Med* 317:588-593, 1987.
- Kuttesch JF, Wexler LH, Marcus RB, et al.: Second malignancies after Ewing's sarcoma: Radiation dose-dependency of secondary sarcomas. *J Clin Oncol* 14:2818-2825, 1996.
- Nicholson HS, Mulvihill JJ, Byrne J: Late effects of therapy in adult survivors of osteosarcoma and Ewing's sarcoma. *Med Pediatr Oncol* 20:6-12, 1992.
- Smith LM, Cox RS, Donaldson SS: Second cancers in long-term survivors of Ewing's sarcoma. *Clin Orthop* 274:275-281, 1992.
- Gasparini M, Lombardi F, Ballerini E, et al.: Long-term outcome of patients with monostotic Ewing's sarcoma treated with combined modality. *Med Pediatr Oncol* 23:406-412, 1994.
- Travis LB, Curtis RE, Hankey BF, et al.: Second cancers in patients with Ewing's sarcoma. *Med Pediatr Oncol* 22:296-297, 1994.
- Zoubek A, Simonitsch I, Panzer-Grümayer ER, et al.: Ewing tumor after treatment of Ki-1+ anaplastic large cell lymphoma. Therapy-associated secondary neoplasm or unrelated coincidence? *Cancer Genet Cytogenet* 83:5-11, 1995.
- Helton KJ, Fletcher BD, Kun LE, et al.: Bone tumors other than osteosarcoma after retinoblastoma. *Cancer* 71:2847-2853, 1993.
- Fisher R, Kaste SC, Parham DM, et al.: Ewing's sarcoma as a second malignant neoplasm in a child previously treated for Wilms' tumor. *J Pediatr Hematol Oncol* 17:76-80, 1995.
- Newton WA Jr, Meadows AT, Shimada H, et al.: Bone sarcomas as second malignant neoplasms following childhood cancer. *Cancer* 67:193-201, 1991.
- Lopez-Andreu JA, Ferris J, Verdeguer A, et al.: Secondary acute promyelocytic leukemia in a child treated with epipodophyllotoxins. *Am J Pediatr Hematol Oncol* 16:384-386, 1994.
- Link MP, Donaldson SS, Kempson RL, et al.: Acute nonlymphocytic leukemia developing during the course of Ewing's sarcoma. *Med Pediatr Oncol* 12:194-200, 1984.